Dr. McCallum-MoviPrep Bowel Preparation Instructions PLEASE READ ALL OF THESE INSTRUCTIONS CAREFULLY

At anytime should you have any questions about the instructions, please contact our office at (520) 327-3454 (during normal business hours) so a staff member can go over them with you.

5 DAYS BEFORE YOUR EXAM

- **STOP** taking all anticoagulants; Coumadin, Heparin, Lovenox, Plavix, Aggranox (if okay with your Primary Care Physician or Cardiologist). If you must continue them, please let us know.
- **STOP** any bulking agents; Metamucil, Citrucel, Gemfibrozil, Ginseng, Ginkgo, Garlic tablets, Vitamin E, Multi Vitamins, & Iron Supplements.

3 DAYS BEFORE YOUR EXAM

- STOP EATING ALL Nuts, Seeds, Red Meat, Leafy Vegetables, Fruit Skins, Salads, Popcorn, Beans (all kinds), Corn, Tomatoes, Salsa, Corn Chips.
- YOU MAY HAVE: Pasta, Eggs, White Rice, Mashed and Sweet Potatoes, Chicken, Fish, White Bread, Cheerios, Pudding, Yogurt, Milk Shakes, Cheese, Creamy Soups, Oatmeal, Bananas, Peeled Apples and Applesauce

DAY BEFORE YOUR EXAM-Clear liquid diet only.

- **Do not** take any oral diabetic medications the day before or the morning of your exam.
- **Insulin dependent diabetics** should take ½ the usual dose of Insulin the day before your exam and ½ the usual dose on the day of your exam. Continue to monitor blood sugar as usual.
- Take any heart and/or blood pressure medication as usual.

Drink as much of the clear liquid diet as you can throughout the day which includes; **Beverages**- Soft drinks (orange, ginger ale, cola, Sprite, 7-Up, etc.), Gatorade, Kool-Aide, strained fruit juices w/o pulp (apple, white grape, lemonade, etc.), water, tea, or coffee. **No** milk or non-dairy creamer. No beverages that are red or purple in color. Soups-low sodium chicken or beef bouillon/broth. Desserts-Hard candies, Jell-O (lemon, Lime or orange; no fruit or toppings), or popsicles (no sherbets or fruit bars).

- Lunch and Dinner-LIQUID DIET ONLY Drink all the clear liquids you like.
- Starting at _____ empty 1 Pouch A and 1 Pouch B into the disposable container, add lukewarm drinking water to the top line of the container. Mix to dissolve.
- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8oz), until the full liter is consumed.
- Drink 16oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep. Continue to drink clear liquids until bedtime.
- Individual responses to laxatives vary. Remain close to a toilet, as multiple bowel movements will occur.

DAY OF YOUR EXAM-Clear liquids only as directed. NO solid food until exam has been completed.

- Starting at_____ empty 1 Pouch A and 1 Pouch B into the disposable container, add lukewarm drinking water to the top line of the container. Mix to dissolve.
- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8oz), until the full liter is consumed.
- Drink 16oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep.
- Take any heart and/or blood pressure medications as usual.
- STOP DRINKING CLEAR LIQUIDS 2 HOURS PRIOR TO YOUR EXAM. Nothing after
- Please bring a list of your medications with you.

YOU MUST HAVE SOMEONE TO DRIVE YOU HOME AFTER YOUR EXAM. WE REQUEST THAT YOUR DRIVER REMAIN DURING YOUR EXAM

Our facility doors open at 6:30am. Should you be scheduled for an early morning procedure, please plan your arrival accordingly.